

МЕЖДУНАРОДНАЯ КОНФЕРЕНЦИЯ

и школа молодых ученых по измерениям, моделированию
и информационным системам для изучения окружающей среды

избранные труды

ENV IRO MIS 20



INTERNATIONAL CONFERENCE

and Early Career Scientists School on Environmental Observations,
Modeling and Information Systems

selected papers

Enviromis 2020 organizers

Institute of Monitoring of Climatic and
Ecological Systems SB RAS

Institute of Numerical Mathematics RAS

Moscow Center for Basic and Applied
Mathematics

Ugra State University

Организаторы Enviromis 2020

Институт мониторинга климатических
и экологических систем СО РАН

Институт вычислительной математики РАН

Московский центр фундаментальной
и прикладной математики

Югорский государственный университет

WCEDAI 2020 organizers

Institute of Monitoring of Climatic
and Ecological Systems SB RAS

Research Computing Center
of Moscow State University

Institute of Numerical Mathematics RAS

Magdeburg-Stendal University
of Applied Sciences

Potsdam Institute
for Climate Impact Research

Организаторы WCEDAI 2020

Институт мониторинга климатических
и экологических систем СО РАН

Научно-исследовательский вычислительный
центр МГУ

Институт вычислительной математики РАН

Магдебург-Стендальский университет
прикладных наук

Потсдамский институт по исследованию
влияния климатических изменений

Thanks

The conference
is supported by RFBR,
grant № 20-05-22040

Благодарности за поддержку

Мероприятие проведено
при финансовой поддержке РФФИ,
проект № 20-05-22040

According to the results of a comparison of baric formations and their characteristics according to electronic bulletins of synoptic maps and maps constructed using reanalysis, it was found that, despite some quantitative differences, at a qualitative level, reanalysis reflects quite well the real picture of their temporal variability.

The comparison of lightning activity registered by different lightning detection systems in the south-east of Western Siberia

^{1,2}Pustovalov K.N., ¹Gorbatenko V.P., ¹Konstantinova D.A., ¹Nechepurenko O.E.

¹Tomsk State University, Tomsk, Russia

²Institute of Monitoring of Climatic and Ecological Systems SB RAS, Tomsk, Russia

E-mail: vpgor@tpu.ru, const.py@yandex.ru

In the second half of XX – early XXI centuries, an increase in the proportion of convective clouds has been noted in Northern Eurasia [1]. Under continued foregoing trend, according to [2], the frequency of Cumulonimbus occurrence and related to them hazardous phenomena will become more frequent. Therefore, an operational identification of convective hazardous phenomena, in particular, thunderstorms, has essential meaning. The aim of this work is to compare the results of lightning registration in the southeast of Western Siberia using different lightning detection systems.

The single-point lightning detector (LD-250) by Boltek installed on account of training laboratory at the Department of Meteorology and Climatology, Geology and Geography Faculty, Tomsk State University (56.46° N, 84.93° E). LD-250 allows detecting lightning discharges within radius of up to 480 km in a fully automatic mode. The data processing carry out in the RimDataLightning author tool, the output file contains information on the polarity and time of the discharge and the coordinates of the lightning strike point [3]. Since the beginning of the XXI century the World Wide Lightning Location Network (WWLLN) is actively elaborating, providing information on the coordinates of discharges over the entire globe through received electromagnetic signals of thunderstorm sources in the VLF band (3–30 kHz) [4].

For comparison, we used the period July 1 to 25, 2019, when numerous thunderstorms were observed over the study area. The coordinates of lightning discharges in one day were mapping with markers superimposed on each other, corresponding to LD-250 and the WWLLN. An analysis of the results showed good qualitative agreement between the positions of the thunderstorm centers. The correlation coefficient between the daily number of lightning discharges detected by LD-250 (the CG, IC and CC lightning) and the WWLLN is 0.89 for the study period. However, the number of lightning discharges by LD-250 is on average 3.3 times higher than captured by the WWLLN.

REFERENCES:

1. Mokhov I.I., Akperov M.G. Tropospheric lapse rate and its relation to surface temperature from re-analysis data // *Izvestiya. Atmospheric and Oceanic Physics*. 2006. V. 42. № 4. P. 430–438.
2. Chernokulsky A.V., Bulygina O.N., Mokhov I. I. Recent variations of cloudiness over Russia from surface daytime observations // *Environmental Research*. 2011. V. 6. № 3. P. 035202.
3. Konstantinova D.A., Gorbatenko V.P. The results of the registration of lightning over the southeastern territory of Western Siberia // *Izvestiya vuzov. Fizika*. 2011. V. 54, № 11/3. P. 156–162 [In Russian].
4. World Wide Lightning Location Network. – URL: <https://wwlln.net/>.

The comparison of lightning activity registered by different lightning detection systems in the south-east of Western Siberia	
Pustovalov K.N., Gorbatenko V.P., Konstantinova D.A., Nechepurenko O.E.	382
Variability of extreme values of bioclimatic indices in the Crimean resorts	
Stefanovich A.A., Voskresenskaya E.N., Lubkov A.S.	383
Changes of extreme air temperature in the Black sea region	
Sukhonos O.Yu., Voskresenskaya E.N.	384
Extreme rates of the temperature according to weather stations in the Northern Hemisphere	
Tartakovsky V.A., Cheredko N.N., Maximov V.G.	385
Influence of extreme weather conditions and snow accumulation regime on ground freezing depth in Russia in winter period 2019/2020	
Frolov D.M.	385
Modelling of extreme concentrations of atmospheric pollutants, their forecast and practical applications in Russia	
Genikhovich E., Kirillova V., Nikolaev V., Yakovleva E., Rumyantsev D., Vylitok N.	386
Mercury emissions from fires in Siberia (case study of the Tomsk Region)	
Lyapina E., Voropay N.	387
Variation of atmospheric instability characteristics against the background of regional surface thermodynamic conditions during fall-winter season	
Martynova Yu., Krupchatnikov V., Gochakov A., Antokhina O.	388
Extreme precipitation and features of their precipitation in the territory of Perm region	
Sivkov B.A.	389
Role of atmospheric circulation in the formation of anomaly warm and anomaly cold winters in the Asian territory of Russia	
Trukhanov A.E., Kochugova E.A.	390
Empiric approach to forecast hot extremes for energy modeling	
Fedotova E.V., Klimenko V.V.	391
Extreme atmospheric precipitation in Western Siberia based on different databases	
Kharyutkina E.V., Loginov S.V., Martynova Yu.V., Kakurkina O.Yu.	392
An empirical method for predicting extreme low winter sea ice extent in the Russian Arctic in the 21st century under global warming (on the example of the Barents Sea)	
Cherenkova E.A., Semenov V.A., Titkova T.B.	392