ТЕЗИСЫ ДОКЛАДОВ

INTERNATIONAL WORKSHOP

**«Multiscale Biomechanics and Tribology
of Inorganic and Organic Systems»**

МЕЖДУНАРОДНАЯ КОНФЕРЕНЦИЯ

**«Перспективные материалы с иерархической структурой
для новых технологий и надежных конструкций»**

**VIII ВСЕРОССИЙСКАЯ НАУЧНО-ПРАКТИЧЕСКАЯ
КОНФЕРЕНЦИЯ С МЕЖДУНАРОДНЫМ УЧАСТИЕМ,
ПОСВЯЩЕННАЯ 50-ЛЕТИЮ ОСНОВАНИЯ
ИНСТИТУТА ХИМИИ НЕФТИ**

«Добыча, подготовка, транспорт нефти и газа»

DOI: 10.17223/9785946218412/1

THE ROLE OF ACIDE-BASE PROPERTIES OF NANOPARTICLES IN CANCER CELLS VIABILITY

¹Lozhkomoev A.S.

¹*Institute of Strength Physics and Materials Science, SB RAS, Tomsk, Russia*

Tumor acidity is caused by increase in protons generation from the anaerobic metabolism of cancer cells through the overexpression of M2-PK, a dimeric isoenzyme of pyruvate kinase [1]. Consequently, tumors have a significantly lower extracellular pH (~6.5–7.1) in comparison to normal tissues (7.4), due to the maintenance of proton efflux pumps [2].

Recent studies have shown that altering the extracellular pH kills tumor cells, reduces cancer metastasis, and reduces the resistance of tumor cells to drugs. This allows us to consider this approach as a promising treatment for cancer, including suboptimal concentrations of chemotherapy drugs [3].

In the present work, the effect of agglomerates of nanosheets of structures based on aluminum hydroxides with different phase composition on the viability of tumor cells was studied.

Studies have shown that the synthesized nano-sheet structures based on oxide and hydroxide phases of aluminum have antitumor properties. At the same time, it was found that γ -Al₂O₃ nanostructures exhibit more pronounced antitumor properties on 4 lines of tumor cells from the five studied (Fig. 1 a).

Fig. 1. Cell viability (a) and isoelectric titration graphs (b) of AlOOH (blue curve), θ -Al₂O₃ (green curve) and γ -Al₂O₃ (red curve)

This may be due to the fact that γ -Al₂O₃ nanostructures adsorb more protons than AlOOH and θ -Al₂O₃. Before reaching pH3.2, an aqueous suspension of synthesized nanostructures with a concentration of 5 mg/ml, 1.95 ml of a 0.01 M HCl solution was required for AlOOH, 2.22 ml for θ -Al₂O₃, and 2.84 ml for γ -Al₂O₃.

1. Sonveaux P., Végran F., Schroeder T., Wergin M. C., Verrax J., Rabbani Z. N., De Saedeleer C. J., Kennedy K. M., Diepart C., Jordan B. F., Kelley M. J., Gallez B., Wahl M. L., Feron O., Dewhirst M. W. Targeting lactate-fueled respiration selectively kills hypoxic tumor cells in mice. // *The Journal of clinical investigation*, 2008. V118. P. 3930–3942.
2. Gillies R., Raghunand N., Garcia-Martin M., Gateby R. pH imaging. // *IEEE Engineering in medicine and biology magazine*, 2004. V. 23. P. 57–64.
4. Walsh M., Fais S., Spugnini E. P., Harguindey S., Izneid T. A., Scacco L., Williams P., Allegrucci C., Rauch C., Omran, Z. Proton pump inhibitors for the treatment of cancer in companion animals. // *Journal of Experimental & Clinical Cancer Research*, 2015. V. 34. P. 93.