

**AB INITIO CALCULATION OF ENERGY LEVELS AND
RO-VIBRATIONAL SPECTRA FOR SiH₄ MOLECULE**

A. V. NIKITIN, *V.E. Zuev Institute of Atmospheric Optics SB RAS, 1, Akademician Zuev square, 634021 Tomsk, Russia*; **Y. S. CHIZHMAKOVA**, *Tomsk State University, 36 Lenin Avenue, 634050 Tomsk, Russian Federation*; **M. REY**, **VI.G. TYUTEREV**, *Groupe de Spectrométrie Moléculaire et Atmosphérique, UMR CNRS 7331, UFR Sciences BP 1039, 51687 Reims Cedex 2, France*

New potential energy surface (PES) and dipole moment surfaces (DMS) of molecule SiH₄ are constructed using extended ab initio CCSD(T) calculations at 19882 nuclear configurations. The PES ¹ and DMS ² analytical representation are determined through an expansion in symmetry adapted products of internal nonlinear coordinates involving 282 and 692 parameters up to the 8th order and 6th order. Lower vibrational and rovibrational levels are calculated ³. Good agreement of calculated fundamentals with observed values was found. The integrated intensities of lower polyads was calculated .

This work is supported by French-Russian LIA SAMIA. Y. S. CHIZHMAKOVA thanks the Tomsk State University and Academic D.I. Mendeleev Fund Program.

¹[doi:10.1016/j.cplett.2010.11.008](https://doi.org/10.1016/j.cplett.2010.11.008), A.V. Nikitin, M. Rey, VI.G. Tyuterev, *Chem. Phys. Lett.*, **501**, 179–186 (2011).

²[doi:10.1016/j.cplett.2013.02.022](https://doi.org/10.1016/j.cplett.2013.02.022), A.V. Nikitin, M. Rey, VI.G. Tyuterev, *Chem. Phys. Lett.*, **565**, 5–11 (2013).

³[doi:10.1039/c3cp50275a](https://doi.org/10.1039/c3cp50275a), M. Rey, A.V. Nikitin, VI.G. Tyuterev, *Phys. Chem. Chem. Phys.*, **15**, 10049–10061 (2013).